

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

31 minutes, 12 seconds. From the second root, $B=78^{\circ}$ 8'=5 hours, 12 minutes, 32 seconds. Length of day=20 hours, 50 minutes, 8 seconds.

Values of B substituted in (1) give for the latitude 36° 44' and 64° 39'.

III. Solution by C. HORNUNG, A. M., Professor of Mathematics, Heidelberg University, Tiffin, O.

Let Z be the zenith, P the pole, N the north point of the horizon, S the place on the horizon where the sun rises, and S' the place of the sun on prime vertical when it has moved half the way to the meridian. Also let ϕ =the required latitude, δ =the sun's declination=23° 27' 15", and P=the hour angle of the sun when rising. Then without allowing for refraction and semi-diameter, we get from the spherical triangles PNS and PZS', $-\cos P$ =tan ϕ tan δ(1); and $\cos \frac{1}{2}P$ = $\cot \phi$ tan δ(2). From (1) and (2) we have $\tan^3 \phi$ -(1/tan δ)tan $^2 \phi$ +2tan δ =0, or $\tan^3 \phi$ -2.31224tan $^2 \phi$ -.18704=0. Whence $\tan \phi$ =2.27614, and ϕ =66° 16′ 54" the required latitude. From (2) P=158° 1′ 28". Hence from sunrise to noon is 10 hours, 32 minutes, 5.8 seconds, and the length of the day is 21 hours, 4 minutes; 11 seconds.

Solved by J. SCHEFFER with result, latitude 36° 43′ 31″, and length of day 14 hours, 31 minutes, 6 seconds.

61. Proposed by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tenn.

The product of n numbers, each the sum of four squares, may be expressed as the sum of four squares in $(48)^{n-1}$ different ways.

Solution by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics, The Russell College, Lebanon, Va.

In Vol. II, No. 2, page 47, it is demonstrated that the product of two numbers each the sum of four squares may be expressed as the sum of four squares in 48 different ways.

Let a_1^2 , a_2^2 , a_3^2 , a_4^2 a_n^2 be the n squares. Then $a_1^2a_2^2 = (m_1^2 + m_2^2 + m_3^2 + m_4^2)$ in 48 ways. $a_1^2a_2^2a_3^2 = (m_1^2a_3^2 + m_2^2a_3^2 + m_3^2a_3^2 + m_4^2a_3^2) = (n_1^2 + n_2^2 + n_3^2 + n_4^2)$ in $4 \times 48 \times 48 = 2^2 \cdot 48^2$ ways. $a_1^2a_2^2a_3^2a_4^2 = (n_1^2a_4^2 + n_2^2a_4^2 + n_3^2a_4^2 + n_4^2a_4^2) = (o_1^2 + o_2^2 + o_3^2 + o_4^2)$ in $4 \times 2^2 \cdot 48^2 \times 48 = 2^4 \cdot 48^3$ ways. $a_1^2a_2^2a_3^2a_4^2a_5^2 = (p_1^2p_2^2p_3^2p_4^2)$ in $4 \times 2^4 \cdot 48^3 \times 48 = 2^6 \cdot 48^4$ ways. $\therefore a_1^2a_2^2a_3^2a_4^2 \dots a_n^2 = (z_1^2 + z_2^2 + z_3^2 + z_4^2)$

in $2^{2n-4} \cdot 48^{n-1}$ ways= $(2)^{2(n-2)} \cdot (48)^{n-1}$ ways.